




Sunflower Water Use Relationships

David C. Nielsen, Research Agronomist, USDA-ARS, Akron, CO

To make an informed decision regarding the use of sunflowers in dryland crop rotations, producers need to know something about the water relationships of this crop in the central Great Plains. At Akron, CO, precipitation during the sunflower growing season averages about 8 inches. But the total amount of water required for sunflower to grow without water stress in this environment averages about 34 in. Some of the 25 in. shortfall in precipitation is made up for by stored soil water. But dryland sunflower production in this area is almost always growing under some water deficit condition which reduces yield from the maximum attainable.

Reports of sunflower water use range widely from 8 to 38 in. per year. The higher values, of course, come from irrigated studies or high rainfall conditions. Sunflower grown where the soil water profile is full will also use higher amounts of water. Sunflower has the ability to extract water from deeper in the soil profile and more from each soil layer than other crops. Average stored soil water use for several crops is shown in Fig. 1. Proso millet extracts about 4 in. from the top 3 ft. of soil, whereas sunflower extracts about 7.5 in. from the 6 ft soil profile. Of course, rotational sequence will make a difference on how much water is available for sunflower to use (Fig. 2). Sunflower following millet in a sunflower-fallow-wheat-millet rotation uses more water than sunflower in a sunflower-fallow-wheat rotation because more water is available following millet than following wheat. At Akron the sunflower soil water use ranges from 3 to 9 in.



A water use/yield function generated from both dryland and irrigated sunflower (Fig. 3) shows that about 150 lb/a of seed is produced for every inch of water use after the first 7 in. of water use. Dryland sunflower yields at Akron have ranged from 100 lb/a to 2000 lb/a. Sunflower yield is very responsive to rainfall received during August and September, just prior to flowering and through seed-filling, (Fig. 4). The relationship for yield vs precipitation (158 lb/a/in) is almost the same as for yield vs total water use (151 lb/a/h).

Using the water use/yield relationship (Fig. 3), the Akron rainfall record from 1964-1997, and assuming 6.5 in. of soil water use, we see a predicted sunflower yield distribution of 570-2080 lb/a, with a median yield of 1320 lb/a (Fig. 5). Nearly half of the years of record would have yields in the 1000-1500 lb/a range.

The very dry condition of the soil profile following sunflower requires significant recharge prior to the planting of the next crop in rotation. It is essential that stalks be left standing in good condition such that snow trapping can occur during storms with strong winds. Tall stalks will trap more snow than short stalks. Overwinter soil water change

is closely related to sunflower silhouette factor (height x diameter x population) due to the increase in snow trapping that occurs with taller stalks at high populations (Fig. 6). A population of 14,000 stalks per acre, one inch in diameter, and 16.7 in. tall would have a silhouette factor of 60, and be expected to recharge the soil water profile by about 4 in. under average winter conditions at Akron.

In summary, precipitation falls way short of evaporative demand in dryland sunflower production in the central Great Plains. Sunflowers are able to extract water from deep in the soil profile and are able to extract more water from each soil layer than other crops. About 150 lb/a of sunflower seed are produced for every inch of water used after the first 7 inches of water use. Based on this water use/yield relationship, sunflower yields should average about 1300 to 1500 lb/a. Because of the very dry condition of the soil following sunflower, it is essential to catch as much snow as possible by leaving tall stalks standing through the winter so that the soil water profile can be recharged.

David C. Nielsen
Research Agronomist
USDA-ARS
Central Great Plains Research Station
P.O. Box 400
Akron, CO 80720

phone: 970-345-0507
fax: 970-345-2088
e-mail: dniesen@lamar.colostate.edu

Visit our website at www.gpsr.colostate.edu/CGPRS

